ZooKeys 1062: 31–47 (2021)
doi: 10.3897/zookeys.1062.64746
https://zookeys.pensoft.net

RESEARCH ARTICLE

A peer-reviewed open-access journal
ZooKeys
Launched to accelerate biodiversity research

New data on the valvatiform-shelled Hydrobiidae (Caenogastropoda, Truncatelloidea) from southern Greece

Sebastian Hofman[1], Jozef Grego[2], Zoltán Fehér[3], Zoltán Péter Erőss[4], Aleksandra Rysiewska[5], Artur Osikowski[6], Andrzej Falniowski[5]

1 *Department of Comparative Anatomy, Institute of Zoology and Biomedical Research, Jagiellonian University, ul. Gronostajowa 9, 30-387 Kraków, Poland* **2** *Horná Mičiná 219, SK-97401 Banská Bystrica, Slovakia* **3** *WWF Hungary H-1141 Álmos vezér útja 69/a, Budapest, Hungary* **4** *Levendula u. 68/4, H-2119, Pécel, Hungary* **5** *Department of Malacology, Institute of Zoology and Biomedical Research, Jagiellonian University, ul. Gronostajowa 9, 30-387 Kraków, Poland* **6** *Department of Animal Reproduction, Anatomy and Genomics, University of Agriculture in Krakow, al. Mickiewicza 24/28, 30-059 Kraków, Poland*

Corresponding author: Sebastian Hofman (s.hofman@uj.edu.pl)

Academic editor: Eike Neubert | Received 21 February 2021 | Accepted 31 August 2021 | Published 12 October 2021

http://zoobank.org/975CEF1A-380F-4DF1-931C-EEBF9A43D984

Citation: Hofman S, Grego J, Fehér Z, Erőss Z-P, Rysiewska A, Osikowski A, Falniowski A (2021) New data on the valvatiform-shelled Hydrobiidae (Caenogastropoda, Truncatelloidea) from southern Greece. ZooKeys 1062: 31–47. https://doi.org/10.3897/zookeys.1062.64746

Abstract

The minute valvatiform-shelled Hydrobiidae are less studied than other hydrobiid gastropods. In this paper, new data on these snails are presented, which have been collected at twelve springs in southern Greece: one in Boeotia, one on Evvoia Island, and ten on the Peloponnese Peninsula. Mitochondrial cytochrome oxidase subunit I (COI) and nuclear histone (H3) have been used to confirm the determinations and infer the relationships of the studied gastropods. They represent the genera *Daphniola*, *Graecoarganiella* and *Isimerope*. New localities, expanding the known geographic ranges, have been presented for *Daphniola hadei* and *Daphniola louisi*. A species of *Daphniola* found at two localities has been identified as a species new to science, and its description, including the shell, penis, and female reproductive organs is given. Possible relationships between *Graecoarganiella* and *Isimerope* are discussed; their representatives are possibly new species. At one locality a single specimen likely represents a new genus: it was found to be most closely related with *Islamia*, but genetically (p-distance) too distant to be congeneric with *Islamia*.

Keywords

COI, H3, molecular phylogeny, new species, Peloponnese, spring gastropods

Copyright *Sebastian Hofman et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Minute freshwater gastropods with depressed trochiform (valvatiform) shells often were classified as belonging to the family Valvatidae Gray, 1840. The first genus described for such hydrobiid snails was *Horatia* Bourguignat, 1887 from Dalmatia (Schütt 1961; Radoman 1965; Szarowska and Falniowski 2014). Hydrobiidae in Greece are still poorly studied, and their microhabitats have drastically disappeared (Szarowska and Falniowski 2011a). The poor knowledge is perhaps of more concern for the valvatiform-shelled hydrobiids, since their low-spired tiny shells are easily overlooked or treated as juveniles. Some authors, for example Schütt (1980), have expanded the ranges of the Central European and North Balkan genera towards Greece, which was criticized by for example, Radoman (1983, 1985). In the present paper, we present the valvatiform-shelled gastropods collected at 12 localities in southern Greece.

Material and methods

The snails were collected by hand or with a sieve at twelve localities listed in Table 1 (Fig. 1) during two trips in 2009 and 2018. Samples were sieved through 500 μm sieve and fixed in 80% analytically pure ethanol, replaced twice, and sorted later. Next, the snails were put in fresh 80% analytically pure ethanol and kept at -20 °C in a refrigerator. The shells were photographed with a Canon EOS 50D digital camera, under a Nikon SMZ18 microscope with dark field. Dissections were performed under a Nikon SMZ18 microscope with dark field, equipped with Nikon DS-5 digital camera. Captured images were used to draw anatomical structures with a graphic tablet. Morphometric parameters of the shell were measured all by the same person using a Nikon DS-5 digital camera and ImageJ image analysis software (Rueden et al. 2017).

Snails for molecular analysis were fixed in 80% ethanol, changed twice, and later stored in 96% ethanol. DNA was extracted from whole specimens; tissues were hydrated in TE buffer (3 × 10 min); then total genomic DNA was extracted with the SHERLOCK extraction kit (A&A Biotechnology), and the final product was dissolved in 20 μl of tris-EDTA (TE) buffer. The extracted DNA was stored at -80 °C at the Department of Malacology, Institute of Zoology and Biomedical Research, Jagiellonian University in Kraków (Poland).

Mitochondrial cytochrome oxidase subunit I (COI), and nuclear histone 3 (H3) loci were sequenced. Details of PCR conditions, primers used and sequencing were given in Szarowska et al. (2016). Sequences were initially aligned in the MUSCLE (Edgar 2004) program in MEGA 7 (Kumar et al. 2016) and then checked in BIOEDIT 7.1.3.0 (Hall 1999). Uncorrected p-distances were calculated in MEGA 7. Estimation of the proportion of invariant sites and the saturation test (Xia 2000; Xia et al. 2003) were performed using DAMBE (Xia 2013). In the phylogenetic analysis, additional sequences from GenBank

Table 1. Geographic coordinates of identified sampling sites, by species. See also the map (Fig. 1). Extraction numbers (in bold) are also given, see also Figures 7, 8.

Id	Site	Coordinates
Daphniola longipenia		
1	W edge of Katarraktis, spring 564 m, Achaia, Peloponnese, Greece; **2A29, 2B24**	38.1014, 21.8328
2	Panagitsa, large spring 500 m, Arcadia, Peloponnese, Greece; **2A32, 2B26, 2B27**	37.7725, 22.2219
Daphniola hadei		
3	3 km W of Megali Vrisi, Laconia, Peloponnese, Greece; **2A27**	37.2267, 22.5222
4	Spring beneath Katafigio Parnonos, Laconia, Peloponnese, Greece; **2B19**	37.2222, 22.6158
5	Kastorio, spring, 3 3 km N of village at aquaducte, Laconia, Peloponnese, Greece; **2B20**	37.1733, 22.2944
Daphniola louisi		
6	Ag. Kiriaki spring N of Kato Kampia, Euboea, Greece; **2A33**	38.5608, 23.8442
Graecoarganiella parnassiana		
7	mouth of Erkinas Gorge, Kria 2, Boeotia, Livadia, Greece **2A28, 2B23**	38.4319, 22.8750
Isimerope semele		
8	Peloponnese, Achaia regional unit, Katarraktis center, spring and limetone cliffs, **2A30**	38.0989, 21.8342
9	Peloponnese, Achaia regional unit, Ag. Georgios (E of Tripotam), Vici spring, **2A31**	37.8525, 21.9397
10	Peloponnese, Achaia regional unit, Ag. Georgios (E of Tripotam), Anastasia spring, **2B21**	37.8517, 21.9408
11	Ladon spring E of Kerasia, Achaia regional unit, 474 m; 462 m alt., Greece, **2A22**	37.8361, 22.1819
cf. *Islamia* sp.		
12	Mili, spring below power station (on the Astros–Argos road), Argolis **2A34**	37.5525, 22.7175



Figure 1. Localities of the sampling sites. For geographic coordinates see Table 1.

were used as references (Table 2). The data were analysed using approaches based on Bayesian Inference (BI) and Maximum Likelihood (ML). We applied the GTR model whose parameters were estimated by RAxML (Stamatakis 2014). In the BI analysis, the GTR + I + Γ model of nucleotide substitution was applied. The model was selected using MrModelTest 2.3 (Nylander 2004). The Bayesian analyses were run using MrBayes v. 3.2.3 (Ronquist et al. 2012) with defaults of most priors. Two simultaneous analyses were performed, each

Table 2. Taxa used for phylogenetic analyses (COI and H3) with their GenBank (GB) accession numbers and references.

Species	COI GB numbers	H3 GB numbers	References
Agrafia wiktori Szarowska & Falniowski, 2011	JF906762	MG543158	Szarowska and Falniowski 2011b; Grego et al. 2017
Alzoniella finalina Giusti & Bodon, 1984	AF367650	-	Wilke et al. 2001
Anagastina zetavalis (Radoman, 1973)	EF070616	-	Szarowska 2006
Belgrandiella kuesteri (Boeters, 1970)	MG551325	MG551366	Osikowski et al. 2018
Dalmatinella fluviatilis Radoman, 1973	KC344541	-	Falniowski and Szarowska 2013
Daphniola dione Radea, Lampri, Bakolitsas & Parmakelis, 2021	MW581160	-	Radea et al. 2021
Daphniola exiqua (A. Schmidt, 1856)	EU047766, JF916470	-	Falniowski et al. 2007; Falniowski and Szarowska 2011a
Dapniola graeca Radoman, 1973	EU047763	-	Falniowski et al. 2007
Daphniola hadei (Gittenberger, 1982)	JF916477, JF916479	-	Falniowski and Szarowska 2011a
Daphniola hadei (Gittenberger, 1982)	MZ093457-MZ093459	MZ265365–MZ265367	present study
Daphniola louisi Falniowski & Szarowska, 2000	EU047769, KM887914, KM887915	-	Falniowski et al. 2007; Szarowska et al. 2014
Daphniola louisi Falniowski & Szarowska, 2000	MZ093456	MZ265364	present study
Daphniola longipenia	MZ093460-MZ093464	MZ265368–MZ265372	present study
Daphniola magdalenae Falniowski, 2015	KT825578, KT825580	-	Falniowski and Sarbu 2015
Ecrobia maritima (Milaschewitsch, 1916)	KX355835	MG551322	Osikowski et al. 2016/Grego et al. 2017
Fissuria boui Boeters, 1981	AF367654	-	Wilke et al. 2001
Graziana alpestris (Frauenfeld, 1863)	AF367641	-	Wilke et al. 2001
Graecoarganiella parnassiana Falniowski & Szarowska, 2011	JN202349, JN202352	-	Falniowski and Szarowska 2011b
Graecoarganiella parnassiana Falniowski & Szarowska, 2011	MZ093454-MZ093455	MZ265362–MZ265363	present study
Grossuana angeltsekovi Glöer & Georgiev, 2009	KU201090	-	Falniowski et al. 2016
Grossuana hohenackeri (Küster, 1853)	KC011749	-	Falniowski et al. 2012
Hauffenia michleri (Kuščer, 1932)	KT236156	KY087878	Falniowski and Szarowska 2015/Rysiewska et al. 2017
Isimerope semele Radea & Parmakelis, 2013	KC841149	-	Radea et al. 2013
Isimerope semele Radea & Parmakelis, 2013	MZ093450-MZ093453	MZ265358–MZ265361	present study
Isimerope sp.	JN202354		Falniowski and Szarowska 2011b
Islamia zermanica (Radoman, 1973)	KU662362	MG551320	Beran et al. 2016; Grego et al. 2017
Islamia sp.	MZ093465	MZ265373	present study
Pontobelgrandiella sp. Radoman, 1978	KU497024	MG551321	Rysiewska et al. 2016/Grego et al. 2017
Radomaniola curta (Küster, 1853)	KC011814	-	Falniowski et al. 2012
Sarajana apfelbecki (Brancsik, 1888)	MN031432	MN031438	Hofman et al. 2019

with 10,000,000 generations, with one cold chain and three heated chains, starting from random trees and sampling the trees every 1000 generations. The first 25% of the trees were discarded as burn-in. The analyses were summarised as a 50% majority-rule tree. Convergence was checked in Tracer v. 1.5 (Rambaut and Drummond 2009). FigTree v. 1.4.4 (Rambaut 2010) was used to visualise the trees. The ML analysis was conducted in RAxML v. 8.2.12 (Stamatakis 2014) using the RAxML-HPC v.8 on XSEDE (8.2.12) tool via the CIPRES Science Gateway (Miller et al. 2010). Bootstrap support was calculated with 1000 replicates and summarised on the best ML tree.

Abbreviations

GNHM Goulandris Natural History Museum, Athens, Greece;
HNHM Hungarian Natural History Museum, Budapest, Hungary;
JG privat collection of Jozef Grego;
ZMUJ Zoological Museum of the Jagiellonian University, Kraków, Poland;
ZPE privat collection of Zoltán Péter Erőss.

Results and discussion

Systematics

Family Hydrobiidae Stimpson, 1865

Genus *Daphniola* Radoman, 1973

Notes. Radoman (1973) described this genus with its type species *D. graeca* Radoman, 1973, from the Daphne Spring in the valley of Tembe, North of Larissa. Schütt (1980) considered *D. graeca* a junior synonym of *Valvata exigua* Schmidt, 1856, described from "Greece". Schütt (1980) designated a neotype from a group of small springs at Agia Paraskevi railway station, situated closely to the Daphne Spring, also in the valley of Tembe in Thessalia, but certainly not close to Thessaloniki as Kabat and Hershler (1993) state. Falniowski and Szarowska (2000) described *Daphniola louisi* from a small spring at the monastery at Kessariani, Athens, Attica. The description was not considered by Bodon et al. (2001), who followed either Schütt (1980) in synonymizing *D. graeca* with *D. exigua*, or Reischütz and Sattmann (1993) in including *Valvata* (*Cincinna*) *hellenica* Westerlund, 1898 in *Daphniola exigua*, thus rendering the genus *Daphniola* monotypic. Falniowski et al. (2007), applying soft-part morphology and anatomy as well as molecular markers, demonstrated the species distinctness of *D. louisi*, and identity of *D. exigua* with *D. graeca*. Gittenberger (1982) described *Horatia hadei*, a new species of *Horatia* he found 5 km SW of Yíthion (Gythion), southern Peloponnese. Later, Falniowski and Szarowska (2011a) collected this gastropod close to the (probably) destroyed type locality, and both, morphology and molecular data confirmed its classification

within the genus *Daphniola*. Radea (2011) described *D. eptalophos* Radea, 2011 from the Parnassos Mountains. However, considering morphology, it is certainly not *Daphniola* especially in the intensively pigmented and massive penis. Its type locality is close (or rather the same) to the type locality of *Graecoarganiella parnassiana* Falniowski & Szarowska, 2011. Thus "*Daphniola etalophos*" is most probably a synonym of *Graecoarganiella parnassiana*, and clearly belongs to *Graecoarganiella* rather than *Daphniola*. Szarowska et al. (2014) found a few juvenile specimens (used for DNA sequencing) of *Daphniola* sp. on each of the two Aegean islands: Rhodos and Khios. Falniowski and Sarbu (2015) described *D. magdalenae* Falniowski, 2015 from the sulphide Melissotripa Cave in Thessalia. Finally, Radea et al. (2021) described *D. dione* Radea, Lampri, Bakolitsas & Parmakelis, 2021 from the Levkas Island (Ionian Sea), using morphology and molecular data. At two localities in North Peloponnese (1 and 2) we found another *Daphniola* taxon, whose morphology and COI sequences do not comply with any known *Daphniola* species.

Daphniola longipenia Grego & Falniowski, sp. nov.

http://zoobank.org/EE503BD0-65ED-4D53-AA37-ED75AB9448DA

Types. Ethanol-fixed specimens, Panagitsa, large spring, Arcadia, Peloponnese, Greece, 37°46'21"N, 22°13'19"E (Fig. 2), altitude 500 m, sieved from sand at the spring head; 26.09.2009; Z.P. Eröss, Z. Fehér, T. Fehér, J. Grego and A. Hunyadi coll., ***holotype***: ZMUJ-M.2137; ***paratypes***: ZMUJ-M.2138-ZMUJ-M.2139, two paratypes; HNHM-105279, 10 wet and 25 dry paratypes, GNHM 39591, 10 paratypes, ZPE 25 dry paratypes, JG F1198, 11 wet and 72 dry paratypes.

GenBank numbers. MZ093460–MZ093464; MZ265368–MZ265372

Diagnosis. Shell minute, valvatiform-trochiform, soft parts slightly pigmented, penis with extremely long and slender filament and small non-glandular outgrowth (lobe) on the left side. Readily distinguished from *D. exigua*, *D. louisi*, *D. magdalenae* and *D. dione* by the proportionally much lower spire of the shell, and the penis with a narrower base and a longer and thinner filament. Differentiated from the geographically (but not molecularly) most close *D. hadei* by the shell with usually lower spire, and the penis with smaller outgrowth and still longer and thinner filament.

Description. Shell (Fig. 3A–E) valvatiform-trochiform, up to 1.00 mm tall, having 3.5 whorls, spire height 10–12% height of shell. Apex flat. Teleoconch whorls moderately convex, evenly rounded, growing rapidly in diameter. Aperture slightly elliptical, parietal lip complete, umbilicus very broad, outer lip simple, orthocline. Teleoconch with delicate growth lines, periostracum pinkish or yellowish. Shell parameters for holotype and a series of paratypes are given in Table 3. Inner and outer sides of operculum smooth. Operculum pinkish. Animal brownish, with some spots of black pigment.

Female reproductive organs (Fig. 4) with a broad loop of the oviduct, a big bursa copulatrix with a long duct, and two moderately small receptacula seminis.

Figure 2. Type locality of *Daphniola longipenia* sp. nov. Panagitsa, Arcadia, Peloponnese **A** spring reservoir **B** spring head.



Figure 3. Shells of gastropods: *D. longipenia* **A–D** locality 2 (holotype, 2A32, 2B26, 2B27) **E** locality 1 (2A29); *Daphniola hadei* **F** locality 3 (2A27) **G** locality 4 (2B19) **H** locality 5 (2B20) **I** *D. louisi*, locality 6 (2A33) **J–K** *Graecoarganiella parnassiana*, locality 7 (2A28, 2B23) **L–O** *Isimerope* **L** locality 8 (2A30) **M** locality 9 (2A31) **N** locality 10 (2B21) **O** locality 11 (2A22) **P** cf. *Islamia* sp., locality 12 (2A34). Scale bar: 1 mm.

Table 3. Shell measurements of *Daphniola longipenia*; specimen symbols as in Figure 3; measured variables: see Figure 6.

	a	*b*	*c*	*d*	*e*	*a*
A – holotype	**0.87**	**0.75**	**0.60**	**0.12**	**0.54**	**121**
B – 2A32	0.88	0.82	0.66	0.10	0.57	127
C – 2B26	0.84	0.72	0.60	0.12	0.48	120
D – 2B27	1.00	0.82	0.66	0.14	0.57	121
E – 2A29	0.74	0.68	0.57	0.10	0.53	127
M	0.866	0.758	0.618	0.116	0.538	123.200
SD	0.093	0.062	0.040	0.017	0.037	3.493
MIN	0.74	0.68	0.57	0.10	0.48	120
MAX	1.00	0.82	0.66	0.14	0.57	127



Figure 4. Pallial and renal section of female reproductive organs of *Daphniola longipenia* [bc – bursa copulatrix, cbc – duct of bursa, ga – albuminoid gland, gn – nidamental gland, gp – gonoporus, ov – oviduct, ovl – loop of (renal) oviduct, rs – seminal receptacles (in black) rs_1 and rs_2 (as defined by Radoman 1973, 1983): rs_1 – distal, rs_2 – proximal]. Scale bar: 250 μm.

Penis (Fig. 5) extremely long and narrow, simple, with an almost vestigial outgrowth proximally on its left edge, and a prominent sharp-terminated filament, vas deferens not visible inside.

Derivatio nominis. The specific epithet *longipenia* refers to the extremely long filament of the penis.

Distribution and habitat. Apart from the type locality (our locality 2), this species was also found in the spring at the W edge of Katarraktis, Achaia, Peloponnese (our locality 1).

Remarks. Westerlund (1898) described *Valvata* (*Cincinna*) *hellenica* Westerlund, 1898, from "Vyteria in Arkadien". Reischütz and Reischütz (2004) identified Westerlund's "Vyteria" as Vitina, situated about 15 km from Panagitsa. They reported *Hauffenia hellenica* (Westerlund, 1898) also from Panagitsa spring. Our *D. longipenia* is most prob-

Figure 5. Penis of *Daphniola longipenia*, bar equals 200 μm.

ably the gastropod reported by them. However, their identification of "Vyteria" as Vitina remains doubtful. The shell of the lectotype of *Valvata hellenica* presented by Reischütz and Sattmann (1993) looks different (Fig. 6) (enormously high and massive body whorl, another size and outline of the aperture). *Valvata hellenica* was reported several times from localities scattered throughout Greece, often in generic combination with *Hauffenia* or *Daphniola*. It can be assumed that these records report more than one species; or it was mentioned as a younger synonym under *Daphniola exigua* (e.g., Bodon et al. 2001). Summarising, the description of a new species is the most appropriate solution.



Figure 6. A shell of lectotype of *Valvata hellenica* presented by Reischütz and Sattmann (1993) **B** shell measurements: a – shell height, b – body whorl breadth, c – aperture height, d – spire height, e – aperture breadth, α – apex angle.

In our trees (Figs 7, 8), *D. longipenia* is clearly distinct from all the other species of *Daphniola* and forms a distinct sister clade opposite to the remaining currently known *Daphniola* species. The high genetic distance (p-distance 0.106) can be found between *D. longipenia* and *D. hadei* (p-distance 0.106), whose localities are most close, and who share the most similar shell morphology, female reproductive organs and penis). In general, the genetic distances between *D. longipenia* and the other *Daphniola* varies from 0.097 (for *D. exigua*) to 0.141 (for *D. magdalenae*) (Table 4).

Daphniola hadei (Gittenberger, 1982)

GenBank numbers. MZ093457–MZ093459; MZ265365–MZ265367

Notes. At the localities 3, 4 and 5 (Fig. 1, Table 1) gastropods were collected, whose shells (Fig. 3F–H), soft parts morphology and anatomy, clearly identified them as belonging to this species. Their molecular data (Figs 7, 8) were identical or nearly identical with the ones published in the GenBank. Their localities are situated somewhat north of the type locality.

Daphniola louisi Falniowski & Szarowska, 2000

GenBank numbers. MZ093456, MZ265364

Notes. The shell morphology (Fig. 3I), soft-part anatomy and molecular data (Figs 7, 8) of the gastropods collected at locality 6 (Fig. 1, Table 1), all showed that they belonged to this species. It has to be noted that the molecular variability in this species (p-distance 0.044) (Fig. 7) is much larger than in *D. hadei* (p-distance 0.013). The new record of *D. louisi*, located on Evvoia Island, considerably expands the range of the species, which so far was only known from Attica. The close phylogenetic relationship with the two juvenile *Daphniola* specimens collected on the Aegean islands, Rhodos and Khios, now combined with the present record from Evvoia, seems to confirm the ideas about the means of dispersal of *Daphniola* from continental Greece to the Aegean islands (Szarowska et al. 2014).

Genus *Graecoarganiella* Falniowski & Szarowska, 2011

Graecoarganiella parnassiana Falniowski & Szarowska, 2011

GenBank numbers. MZ093454–MZ093455; MZ265362–MZ265363

Notes. Falniowski and Szarowska (2011b) described a new, so far monotypic, genus of Hydrobiidae from Greece, Parnassus Mountains, S of Eptalofos, N of Kalania, found in a cistern and a small spring in a grassy pasture on a mountain pass. The type species, *G. parnassiana* Falniowski & Szarowska, 2011, is so far known only from the type locality. At the locality 7 (Fig. 1, Table 1), mouth of Erkinas Gorge, Kria 2,



Figure 7. Phylogenetic tree for COI showing relationships between the studied snails. Bootstrap supports (>60%) and Bayesian probabilities are given.



Figure 8. Phylogenetic tree inferred from connected COI and H3 sequences. Bootstrap supports (>60%) and Bayesian probabilities are given.

Table 4. P-distances for COI between main clades of the *Daphniola*.

	D. longipenia	***D. hadei***	***D. dione***	***D. exiqua***	***D. magdalenae***
D. longipenia	–				
D. hadei	0.105	–			
D. dione	0.088	0.080	–		
D. exiqua	0.097	0.092	0.087	–	
D. magdalenae	0.141	0.153	0.133	0.154	–
D. louisi	0.121	0.103	0.097	0.110	0.122

Boeotia, Livadia, we found gastropods, whose shells (Fig. 3J–K), and soft-part morphology indicated they belonged to *Graecoarganiella*, and were practically identical to *G. parnassiana*. Anatomy was not studied since the material was scarce and not fixed well enough. Our locality 7 is not far (about 35 km) from the type locality of *G. parnassiana*. The molecular data – partial sequences of COI – of our population showed

rather high distinctness (Fig. 7). However, as can be seen in the same phylogram, these differences (p-distance 0.038) are a little lower than the ones within *Daphniola louisi* (0.044). Thus, inclusion of our new population in *Graecoarganiella parnassiana* is seemingly justified.

Genus *Isimerope* Radea & Parmakelis, 2013

Isimerope semele Radea & Parmakelis, 2013

GenBank numbers. MZ093450–MZ093453; MZ265358–MZ265361

Notes. When describing *Graecoarganiella parnassiana* from the Parnassus Mts., Falniowski and Szarowska (2011b) reported three young hydrobiid specimens found at Mainalo Mountain, Peloponnese, WSW of Piana, WNW of Tripolis, in a medium-sized spring and cistern. Their COI sequence was interpreted as indicating a distinct species congeneric with *Graecoarganiella parnassiana*. Later, Radea et al. (2013) found other species at Megali Vrisi, Pharmakas Mt., and described it as a representative of a new monotypic genus *Isimerope*, with *I. semele* as the type species. In our tree (Fig. 7) *Graecoarganiella* and *Isimerope* are quite distinct (p-distance 0.096), but form a well-supported clade (bootstrap value of 100%, Bayesian probability 1.0). The shells are very similar, and the same holds true for the radulae. The lack of a ctenidium, and egg capsules laid in the umbilicus of the shell, might be considered as unique shared character states. The penes and female reproductive organs of the compared taxa do not differ more than could be expected by different season of collection or fixation technique.

At the four localities: 8, 9, 10 and 11 (Fig. 1, Table 1) we collected gastropods, whose shells (Fig. 3L–O), soft parts morphology (not well-fixed material reduced the possible examination) and molecular data (Fig. 7) showed them as belonging to *Isimerope*. Again, as in the case of *Graecoarganiella*, our specimens of *Isimerope* may represent distinct species, but as in *Daphniola*, the molecular differences may be considered as within- species level variation (p-distance 0.035).

Our molecular data clearly show the close relationship of *Isimerope* and *Graecoarganiella*, contradicting their classification to different subfamilies (Belgrandiinae de Stefani, 1877 and Horatiinae D. W. Taylor, 1966, respectively), as stated in WORMS (WoRMS Editorial Board 2021). Both more anatomical and molecular data, as well as a broad-scale revision of the systematics of the Truncatelloidea proposed by Bouchet et al. (2017) are badly needed.

cf. *Islamia* sp.

GenBank numbers. MZ093465; MZ265373

Notes. At the locality 12, in Mili, Argolis, in a spring below the power station, a gastropod was found (Fig. 3P), whose molecularly inferred phylogenetic position (Fig. 8) remains enigmatic. Its sister taxon is *Islamia* Radoman, 1973. The clade's bootstrap sup-

port for two concatenated loci is 100%, strongly suggesting that both mOTUs belong to the same taxon, but the genetic distance between them seems too high (p-distance 0.135). The p-distances in COI were 0.109 and 0.138 between this taxon and *Fissuria* Boeters, 1981 and *Alzoniella* Giusti & Bodon, 1984, respectively, although the shell morphology still suggests an affiliation with *Islamia*. Anyway, with only one more shell and lack of molecular data on the other Greek *Islamia* species, a justified taxonomic decision has to be postponed until more, and better, preserved specimens are available.

Acknowledgements

We would like to thank to András Hunyady (Budapest, Hungary) and Tamás Fehér (Budapest, Hungary) for the assistance during the 2018 field trip. The study was supported by a grant from the National Science Centre 2017/25/B/NZ8/01372 to Andrzej Falniowski.

References

Arconada B, Ramos M-A (2006) Revision of the genus *Islamia* Radoman, 1973 (Gastropoda, Caenogastropoda, Hydrobiidae) on the Iberian Peninsula and description of two new genera and three new species. Malacologia 48: 77–13.

Beran L, Bodon M, Cianfanelli S (2014) Revision of "*Hauffenia jadertina*" Kuščer 1933, and description of a new species from Pag island, Croatia (Gastropoda: Hydrobiidae). Journal of Conchology 41: 585–601.

Beran L, Osikowski A, Hofman S, Falniowski A (2016) *Islamia zermanica* (Radoman, 1973) (Caenogastropoda: Hydrobiidae): morphological and molecular distinctness. Folia Malacologica 24: 25–30. https://doi.org/10.12657/folmal.024.004

Bodon M, Cianfanelli S (1996) A new valvatoid-shelled stygobiont hydrobiid from Slovenia (Gastropoda: Prosobranchia: Hydrobiidae). Basteria 60: 19–26.

Bodon M, Manganelli G, Giusti F (2001) A survey of the European valvatiform hydrobiid genera, with special reference to *Hauffenia* Pollonera, 1898 (Gastropoda: Hydrobiidae). Malacologia 43: 103–215.

Bouchet P, Rocroi J-P, Hausdorf B, Kaim A, Kano Y, Nutzel A, Parkhaev P, Schrödl M, Strong EE (2017) Revised classification, nomenclator and typification of gastropod and monoplacophoran families. Malacologia 61: 1–526. https://doi.org/10.4002/040.061.0201

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research 32: 1792–1797. https://doi.org/10.1093/nar/gkh340

Falniowski A, Sarbu S (2015) Two new Truncatelloidea species from Melissotrypa Cave in Greece (Caenogastropoda). ZooKeys 530: 1–14. https://doi.org/10.3897/zookeys.530.6137

Falniowski A, Szarowska M (2000) A new species of *Daphniola* Radoman, 1973 (Gastropoda: Hydrobiidae) from Greece. Folia Malacologica 8: 181–188. https://doi.org/10.12657/folmal.008.013

Falniowski A, Szarowska M (2011a) Genus *Daphniola* Radoman, 1973 (Caenogastropoda: Hydrobiidae) in the Peloponnese, Greece. Folia Malacologica 19: 131–137. https://doi.org/10.2478/v10125-011-0020-9

Falniowski A, Szarowska M (2011b) A new genus and new species of valvatiform hydrobiid (Rissooidea; Caenogastropoda) from Greece. Molluscan Research 31: 189–199.

Falniowski A, Szarowska M (2013) Phylogenetic relationships of *Dalmatinella fluviatilis* Radoman, 1973 (Caenogastropoda: Rissooidea). Folia Malacologica 21: 1–7. https://doi.org/10.12657/folmal.021.001

Falniowski A, Szarowska M (2015) Species distinctness of *Hauffenia michleri* (Kuščer, 1932) (Caenogastropoda: Truncatelloidea: Hydrobiidae). Folia Malacologica 23: 193–195. https://doi.org/10.12657/folmal.023.016

Falniowski A, Szarowska M, Grzmil P (2007) *Daphniola* Radoman, 1973 (Gastropoda: Hydrobiidae): shell biometry, mtDNA, and the Pliocene flooding. Journal of Natural History 41: 2301–2311. https://doi.org/10.1080/00222930701630733

Falniowski A, Szarowska M, Glöer P, Pešić V (2012) Molecules vs morphology in the taxonomy of the *Radomaniola/Grossuana* group of Balkan Rissooidea (Mollusca: Caenogastropoda). Journal of Conchology 41: 19–36.

Falniowski A, Georgiev D, Osikowski A, Hofman S (2016) Radiation of *Grossuana* Radoman, 1973 (Caenogastropoda: Truncatelloidea) in the Balkans. Journal of Molluscan Studies 82: 305–313. https://doi.org/10.1093/mollus/eyv062

Gittenberger E (1982) Eine neue *Horatia* art aus Griechenland (Prosobranchia, Hydrobiidae). Basteria 46: e68.

Grego J, Hofman S, Mumladze L, Falniowski A (2017) *Agrafia* Szarowska & Falniowski, 2011 (Caenogastropoda: Hydrobiidae) in the Caucasus. Folia Malacologica 25: 237–247. https://doi.org/10.12657/folmal.025.025

Haase M (1993) *Hauffenia kerschneri* (Zimmermann 1930): zwei Arten zweier Gattungen (Caenogastropoda: Hydrobiidae). Archiv für Molluskenkunde 121: 91–109. https://doi.org/10.1127/arch.moll/121/1992/91

Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95–98.

Hofman S, Osikowski A, Rysiewska A, Grego J, Glöer P, Dmitrović D, Falniowski A (2019) *Sarajana* Radoman, 1975 (Caenogastropoda: Truncatelloidea): premature invalidation of a genus. Journal of Conchology 43: 407–418.

Kabat AR, Hershler R (1993) The prosobranch snail family Hydrobiidae (Gastropoda: Rissooidea): review of classification and supraspecfic taxa. Smithsonian Contributions to Zoology 547: 1–94. https://doi.org/10.5479/si.00810282.547

Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. Molecular Biology and Evolution 33: 1870–1874. https://doi.org/10.1093/molbev/msw054

Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Proceedings of the Gateway Computing Environments Workshop (GCE), 14 Nov, New Orleans, LA, 1–8. https://doi.org/10.1109/GCE.2010.5676129

Nylander JAA (2004) MrModeltest v.2. Program distributed by the author. Uppsala: Evolutionary Biology Centre, Uppsala University.

Osikowski A, Hofman S, Georgiev D, Kalcheva S, Falniowski A (2016) Aquatic snails *Ecrobia maritima* (Milaschewitsch, 1916) and *E. ventrosa* (Montagu, 1803) (Caenogastropoda: Hydrobiidae) in the east Mediterranean and Black Sea. Annales Zoologici 66: 477–486. https://doi.org/10.3161/00034541ANZ2016.66.3.012

Radea C (2011) A new species of hydrobiid snails (Mollusca, Gastropoda, Hydrobiidae) from central Greece. ZooKeys 138: 53–64. https://doi.org/10.3897/zookeys.138.1927

Radea C, Lampri PN, Bakolitsas K, Parmakelis A (2021) A new hydrobiid species (Caenogastropoda, Truncatelloidea) from insular Greece. Zoosystematics and Evolution 97: 111–119. https://doi.org/10.3897/zse.97.60254

Radea C, Parmakelis A, Mourikis T, Triantis KA (2013) *Isimerope*, a new genus of Hydrobiidae (Caenogastropoda: Rissooidea) from Greece. Journal of Molluscan Studies 79: 168–176. doi:10.1093/mollus/eyt010

Radoman P (1965) Speciation der Gattung *Horatia* im Flußtal der Cetina. Archiv für Molluskenkunde 95: 243–253.

Radoman P (1973) New classification of fresh and brackish water Prosobranchia from the Balkans and Asia Minor. Posebna Izdanja, Prirodne Museum Beograd 32: 1–30.

Radoman P (1983) Hydrobioidea a superfamily of Prosobranchia (Gastropoda). I. Systematics. Monographs Serbian Academy of Sciences and Arts, DXLVII, Department Sciences 57: 1–256.

Radoman P (1985) Hydrobioidea, a superfamily of Prosobranchia (Gastropoda). II. Origin, zoogeography, evolution in the Balkans and Asia Minor. Faculty of Science – Department of Biology Monographs, 1, Institute of Zoology Beograd 1: 1–173.

Rambaut A (2010) FigTree v1.3.1. http://tree.bio.ed.ac.uk/software/figtree

Rambaut A, Drummond AJ (2009) Tracer v1.5. http://beast.bio.ed.ac.uk/Tracer

Reischütz PL, Sattmann H (1993) Beiträge zur Nomenklatur der europäischen Binnenmollusken, V. Die Taxa der Hydrobioidea des griechischen Festlands mit valvatoiden Gehäuse und Festlegung eines Lectotypus von *Valvata* (*Cincinna*) *hellenica* Westerlund, 1898 (Gastropoda: Prosobranchia). Heldia 2: 51–52.

Reischütz A, Reischütz PL (2004) Hellenikä pantoia, 7: *Hauffenia edlingeri* nov. spec. (Gastropoda: Prosobranchia: Hydrobiidae) aus Arkadien (Peloponnes, Griechenland). Nachrichtenblatt der Ersten Vorarlberger Malakologischen Gesellschaft 12: 1–2.

Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) Efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61: 539–542. https://doi.org/10.1093/sysbio/sys029

Rueden DT, Schindelin J, Hiner MC, Dezonia BE, Walter AE, Arena ET, Eliceiri KW (2017) ImageJ2: ImageJ for the next generation of scientific image data. BMC Bioinformatics 18: 529. https://doi.org/10.1186/s12859-017-1934-z

Rysiewska A, Georgiev D, Osikowski A, Hofman S, Falniowski A (2016) *Pontobelgrandiella* Radoman, 1973 (Caenogastropoda: Hydrobiidae): A recent invader of subterranean waters? Journal of Conchology 42: 193–203.

Rysiewska A, Prevorčnik S, Osikowski A, Hofman S, Beran L, Falniowski A (2017) Phylogenetic relationships in *Kerkia* and introgression between *Hauffenia* and *Kerkia* (Caenogas-

tropoda: Hydrobiidae). Journal of Zoological Systematics and Evolutionary Research 55: 106–117. https://doi.org/10.1111/jzs.12159

Schütt H (1961) Das Genus *Horatia* Bourguignat. Archiv für Molluskenkunde 90: 69–77.

Schütt H (1980) Zur Kenntnis griechischer Hydrobiiden. Archiv für Molluskenkunde 110: 115–149.

Stamatakis A (2014) RaxML Version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30: 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Szarowska M (2006) Molecular phylogeny, systematics and morphological character evolution in the Balkan Rissooidea (Caenogastropoda). Folia Malacologica 14: 99–168. https://doi.org/10.12657/folmal.014.014

Szarowska M, Falniowski A (2011a) Destroyed and threatened localities of rissooid snails (Gastropoda: Rissooidea) in Greece. Folia Malacologica 19: 35–39. https://doi.org/10.12657/folmal.028.025

Szarowska M, Falniowski A (2011b) An unusual, flagellum-bearing hydrobiid snail (Gastropoda, Rissooidea, Hydrobiidae) from Greece, with descriptions of a new genus and a new species. Journal of Natural History 45: 2231–2246. https://doi.org/10.1080/00222933.2011.591067

Szarowska M, Falniowski A (2014) *Horatia* Bourguignat, 1887: is this genus really phylogenetically very close to *Radomaniola* Szarowska, 2006 (Caenogastropoda: Truncatelloidea)? Folia Malacologica 22: 31–39. https://doi.org/10.12657/folmal.022.003

Szarowska M, Hofman S, Osikowski A, Falniowski A (2014) *Daphniola* Radoman, 1973 (Caenogastropoda: Truncatelloidea) at east Aegean islands. Folia Malacologica 22: 11–20. https://doi.org/10.12657/folmal.022.021

Szarowska M, Osikowski A, Hofman S, Falniowski A (2016) *Pseudamnicola* Paulucci, 1878 (Caenogastropoda: Truncatelloidea) from the Aegean Islands: a long or short story? Organisms Diversity and Evolution 16: 121–139. https://doi.org/10.1007/s13127-015-0235-5

Westerlund CA (1898) Novum specilegium malacologicum. Neue Binnenconchylien aus der paläarktischen Region. - Ezhegodnik, Zoologicheskago Muzeya Imperatorskoi Akademii Nauk [Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St.-Pétersbourg] 3(2): 155–183.

Wilke T, Davis GM, Falniowski A, Giusti F, Bodon M, Szarowska M (2001) Molecular systematics of Hydrobiidae (Mollusca: Gastropoda: Rissooidea): testing monophyly and phylogenetic relationships. Proceedings of the Academy of Natural Sciences of Philadelphia 151: 1–21. https://doi.org/10.1635/0097-3157(2001)151[0001:MSOHMG]2.0.CO;2

WoRMS Editorial Board (2021) World Register of Marine Species. http://www.marinespecies.org. [Accessed at VLIZ 2021-08-30]

Xia X (2000) Data analysis in molecular biology and evolution. Kluwer Academic Publishers, Boston, Dordrecht & London.

Xia X (2013) DAMBE: A comprehensive software package for data analysis in molecular biology and evolution. Molecular Biology and Evolution 30: 1720–1728. https://doi.org/10.1093/molbev/mst064

Xia X, Xie Z, Salemi M, Chen L, Wang Y (2003) An index of substitution saturation and its application. Molecular Phylogenetics and Evolution 26: 1–7. https://doi.org/10.1016/S1055-7903(02)00326-3